Meningococcal Disease





Meningococcal infection (or meningococcal disease) is caused by a bacteria called meningococcus. Meningococcal infection is uncommon, and not easily spread, but it can cause serious complications, including:

- Meningitis an infection of the covering of the brain and spinal cord
- Septicaemia an infection of the blood
- Infections in other parts of the body, such as the joints.

Meningococcal infections can start suddenly and become extremely serious very quickly. They can cause life-long disabilities, or death in about one in 10 people affected.

Immunisation against meningococcal bacteria is the best protection against meningococcal disease.

Meningococcal ACWY Vaccine

Meningococcal ACWY vaccine is provided as a single dose, free through the National Immunisation Program for:

- Children at 12 months of age
- Adolescents aged approximately 14 to 16 years through school based immunisation programs and those aged 15-19 as part of an ongoing catch up program
- People with particular medical risk conditions

Meningococcal B (Bexsero Vaccine)

Moira Shire Council Immunisation Service can provide Meningococcal B vaccine at a cost of \$155 per dose without a prescription.

- Can be given to babies from 6 weeks of age. The number of doses required varies with age
- Aboriginal and Torres Strait Islander children under 2 are funded to receive this vaccine through the National Immunisation Program

Paracetamol

Paracetamol is recommended 30 minutes before, or as soon as practicable after meningococcal B vaccine administration, for children younger than two years of age. Two further doses of paracetamol are recommended six hours apart, regardless of whether fever is present.

For further information on meningococcal disease and immunisation: https://www.health.gov.au/health-topics/immunisation/immunisation-services/meningococcal-immunisation-service-0

For inquiries or to schedule an appointment contact Moira Shire Council Immunisation Team on 5871 9222.